

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁵ : C12P 21/08, C12N 5/20 G01N 33/574, 33/577, A61K 39/395	A1	(11) International Publication Number: WO 93/25700 (43) International Publication Date: 23 December 1993 (23.12.93)
(21) International Application Number: PCT/EP93/01533 (22) International Filing Date: 16 June 1993 (16.06.93) (30) Priority data: RM92A000457 17 June 1992 (17.06.92) IT (71) Applicant (for all designated States except US): ISTITUTO SUPERIORE DI SANITA' [IT/IT]; Viale Regina Elena, 299, I-00161 Rome (IT). (72) Inventor; and (75) Inventor/Applicant (for US only) : CIANFRIGLIA, Maurizio [IT/IT]; Viale Mencacci, 3, I-00042 Anzio (IT). (74) Agent: GRUBB, Philip, W.; B.A. Yorke & Co., Coomb House, 7 St. John's Road, Isleworth, Middlesex TW7 6NH (GB).		(81) Designated States: AU, BB, BG, BR, CA, CZ, FI, HU, JP, KP, KR, LK, MG, MN, MW, NO, NZ, PL, RO, RU, SD, SK, UA, US, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i>
(54) Title: MONOCLONAL ANTIBODIES TO GLYCOPROTEIN P (57) Abstract <p>The invention provides a monoclonal antibody that recognises a structurally continuous and extracellularly-located epitope of human glycoprotein P (Pgp) which consists of a specific continuous epitope-forming amino acid sequence which is preferably located on the fourth extracellular loop of human Pgp and especially contained within the peptide of SEQUENCE ID NO. 1. Preferred antibodies according to the invention bind human Pgp specifically and with high avidity and thus may be used to identify human MDR cells when present as only a very low proportion of a cell population or when expressing human Pgp at only very low levels. The monoclonal antibody may be a whole antibody or antigen binding fragment thereof and may be prepared by hybridoma or recombinant DNA techniques. The monoclonal antibody is useful for the identification or purification of cells which express human Pgp, e.g. when contained in heterogeneous cell populations, and for monitoring the multi-drug resistant status of cells, e.g. tumour cells.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	FR	France	MR	Mauritania
AU	Australia	GA	Gabon	MW	Malawi
BB	Barbados	GB	United Kingdom	NL	Netherlands
BE	Belgium	GN	Guinea	NO	Norway
BF	Burkina Faso	GR	Greece	NZ	New Zealand
BC	Bulgaria	HU	Hungary	PL	Poland
BJ	Benin	IE	Ireland	PT	Portugal
BR	Brazil	IT	Italy	RO	Romania
CA	Canada	JP	Japan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SK	Slovak Republic
CI	Côte d'Ivoire	LJ	Liechtenstein	SN	Senegal
CM	Cameroon	LK	Sri Lanka	SU	Soviet Union
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	MC	Monaco	TG	Togo
DE	Germany	MG	Madagascar	UA	Ukraine
DK	Denmark	ML	Mali	US	United States of America
ES	Spain	MN	Mongolia	VN	Viet Nam
FI	Finland				

Monoclonal Antibodies to glycoprotein P

The present invention relates to monoclonal antibodies which have specificity for P-glycoprotein (Pgp). More particularly, the invention relates to monoclonal antibodies to an extracellular epitope of human Pgp and to their preparation and diagnostic and clinical uses.

The present description refers to various publications by reference numbers in brackets and these are listed towards the end of the text.

The development by Kohler and Milstein (1) of somatic cell fusion techniques for the production of hybrid cell lines which produce monoclonal antibodies has made it possible to prepare unlimited quantities of homogeneous antibody products which have defined specificity for single antigenic determinants. The portion of a protein antigenic determinant that is recognised by and reacts with an antibody is known, and referred to herein, as the epitope and consists of one or more specific epitope-forming amino acid sequences.

Glycoprotein P is a plasma membrane protein which is over-expressed on the surfaces of cell lines or human tumour cells that exhibit an intrinsic or acquired multidrug-resistant (MDR) phenotype. The expression of this protein gives the cells the ability to multiply actively in the presence of concentrations of cytostatic drugs that are toxic for non-MDR parent cells. Transfection studies using cloned Pgp cDNAs have demonstrated the direct role of this protein in mediating the MDR phenotype (2). Pgp is thought to function as an energy-dependent drug efflux pump.

The amino acid sequence of Pgp has been deduced from the nucleotide sequences of the genes that confer the MDR phenotype in recipient cells (2, 3, 4). The secondary structure of Pgp has been predicted from analysis of its primary structure and is that

of a transmembranal protein consisting of 12 hydrophobic transmembranal helices linked via 6 hydrophilic extracellular loops and 2 large cytoplasmic domains encoding 2 ATP-binding sequences.

A number of different monoclonal antibodies to Pgp have been isolated and characterised, and the specificities of some of these are reviewed by Cenciarelli et al. (5). More recently further monoclonals have been described which have specificity for the extracellular domain of Pgp (6, 7, 8). However monoclonal antibodies to Pgp for which the specific epitope-forming amino acid sequences have been characterised react either with the cytoplasmic domain of Pgp or with extracellular epitopes comprising discontinuous portions of the amino acid sequence of the protein. For example, Georges et al. (8) have mapped the epitope of the monoclonal antibody MRK-16, described by Hamada and Tsuruo (9), which comprises peptides present on 2 of the 6 extracellular peptide loops (the 1st and the 4th), predictable from the primary amino acid sequence.

Monoclonal antibodies to Pgp have been proposed for diagnostic and clinical uses, e.g. to monitor the level of Pgp expression on, and thereby the multi drug resistance (mdr) status of, human cells. Also such antibodies may find uses in the inhibition of Pgp-mediated MDR (7).

None of the known monoclonal antibodies to Pgp recognises a human-specific, extracellular epitope of glycoprotein P consisting of a continuous epitope-forming amino acid sequence. Moreover there is a continuing need for antibodies which are capable of recognising low levels of Pgp; for instance, when expressed on MDR cells at low levels or when the MDR cells are present as only a low proportion of a cell population. Monoclonal antibodies capable of recognising such human-specific epitopes and low Pgp levels would be useful for immunological testing.

Using the above mentioned somatic cell fusion techniques, a hybridoma secreting a monoclonal antibody (MM4.17) has been obtained which is capable of selective, high affinity binding to a previously unknown continuous extracellular epitope of human Pgp .

Accordingly the present invention provides a monoclonal antibody that recognises a structurally continuous and extracellularly-located epitope of human Pgp consisting of a continuous epitope-forming amino acid sequence.

For the purposes of the present description "human Pgp" refers to the product of the human MDR1 gene. Also the term "monoclonal antibody" encompasses whole antibody molecules and antigen binding fragments thereof, including FAb, F(Ab')₂, Fv fragments and single domain antibodies, and is not limited to products derived from hybridoma cell lines but also includes antibody molecules and antigen-binding fragments thereof when produced by recombinant DNA techniques, i.e. expressed from cloned antibody coding DNA sequences.

The epitope-forming amino acid sequence which is recognised by the antibody may be located on any one of the extracellular loops of human Pgp. Preferably the epitope-forming amino acid sequence is located on the fourth loop of human Pgp. More preferably the epitope is contained in the human Pgp peptide having the sequence: FTRIDDPETKRQNSNL (SEQUENCE ID NO.1 of the accompanying SEQUENCE LISTING). Even more preferably, the epitope comprises at least 5 contiguous amino acid residues of the 8 amino acids of the peptide consisting of residues 2 to 9 inclusive of SEQUENCE ID NO.1 and still more preferably the 5 amino acids are amino acid residues 3 to 7 inclusive of SEQUENCE ID NO.1.

In a particular embodiment of the invention the monoclonal antibody may have a variable domain amino acid sequence comprising Complementarity Determining Regions (CDRs) in which each CDR is the same as or a variant of the corresponding CDR

given hereinafter in SEQUENCE ID NOs. 2 and 3 (i.e. the CDRs of the heavy chain variable domain of the monoclonal antibody MM4.17).

In the amino acid sequence given in SEQUENCE ID NOs. 2 and 3 CDR1 corresponds to amino acid residues 32 to 36 inclusive, CDR2 to amino acid residues 51 to 67 inclusive and CDR3 to amino acid residues 100 to 111 inclusive. The remainder of the amino acid sequence given in SEQUENCE ID NOs. 2 and 3 comprises framework regions (FR1, FR2, FR3 and FR4 in N terminal to C terminal direction) which may be changed for other framework regions without substantially effecting the antigen binding characteristics of the variable domain.

For the purposes of the present description a variant of a CDR sequence is a sequence which is at least 70%, preferably at least 80%, more preferably at least 90%, particularly at least 95% homologous to a reference CDR sequence, e.g. CDR1, CDR2 or CDR3 of the amino acid sequence given in SEQUENCE ID NOs. 2 and 3.

Surprisingly, compared with the known monoclonal antibodies, the monoclonal antibody MM4.17 has a very high degree of affinity for MDR cells and is capable of detecting even the minimal levels of acquired multidrug resistance of CEM-VBL10 cells (10, 11). MM4.17 is also capable of detecting the acquired multidrug resistance of CEM-VBL16 cells (11). The relative resistance (RR) of this cell line to vinblastine is 6.25. The CEM-VBL10 and CEM-VBL16 cell lines are MDR derivatives of the drug sensitive CEM cell line (see reference 11, deposited as ATCC CCL 119) obtained by selection in medium containing 10 ng/ml and 16 ng/ml respectively of vinblastine.

Thus in a particularly preferred embodiment the invention includes monoclonal antibodies of the invention which have binding affinities for Pgp similar to that of MM4.17, e.g. monoclonal antibodies capable of detecting the acquired multidrug resistance of CEM-VBL10 or CEM-VBL16 cells or similar MDR cells.

A comparison of the binding affinities of the MM4.17 antibody and previous anti-Pgp monoclonals C219 (15), C494 (15), JSB1 (16), MRK-16 (9) and MC57 (5) for Pgp on drug-sensitive parent CEM and MDR derivative cell lines CEM-VBL10 and CEM-VBL100 is given hereinafter in Example 6. The MDR cell lines CEM-VBL10, CEM-VBL16 and CEM-VBL100 are described by and obtainable from Beck et al. (11). Similar MDR cell lines may be obtained substantially as described by Beck et al..

Monoclonal antibodies according to the invention may be prepared using standard techniques. Thus hybridoma cell lines may be prepared by somatic cell fusion techniques using spleen cells obtained from animals which have been immunised with human MDR cells. Any suitable human MDR cell line may be used including MDR cell lines derived from drug sensitive parent cell lines, such as the MDR cell line CEM-VBL100 described by Beck et al. (11) or similar MDR cell lines. CEM-VBL100 is an MDR derivative of the drug sensitive CEM cell line which is deposited with the American type culture Collection as ATCC CCL 119.

More conveniently peptides corresponding to extracellular domains of human Pgp or fragments thereof may be used for immunisation. Any suitable human Pgp extracellular domain peptide may be used. For example, peptides corresponding to the fourth loop of Pgp or fragments thereof, such as the peptide of SEQUENCE ID NO.1 or the fragments thereof discussed above may be used as antigens.

Preferably immunisation procedures which favour the generation of high affinity monoclonals are used, such as the procedures described by Cianfriglia et al. (12), and similar procedures.

Hybridomas which produce the desired monoclonal antibodies may be identified by their ability to bind preferentially to human MDR cells as compared to drug-sensitive cells, followed by selection for those which are specific for structurally continuous extracellularly-located epitopes. Preferably peptides corresponding to structurally continuous extracellular domains

of human Pgp or parts thereof, such as the preferred peptides discussed above, are used to screen for the desired monoclonal antibodies.

Monoclonal antibodies may be prepared from the hybridoma cell lines using cell culture and ascites techniques. Antibody fragments may be prepared using appropriate techniques such as enzymic cleavage. Recombinant products may be produced by processes involving transformation or transfection of suitable host cells with DNA sequences which code for the whole molecule or fragment monoclonal products and culturing of the transformed host cells to produce the products.

The antibody can be used advantageously: to detect the presence of Pgp in normal and tumour tissues with the aid of immunofluorescent or immunochemical methods; in in-vitro tests to identify cells with the MDR phenotype; for methods of enriching specific cell populations in which Pgp is present (stem cells, cells infected with HIV, T lymphocytes).

Thus the invention also includes suitably labelled, e.g. fluorescently labelled monoclonal antibodies according to the invention, as well as products in forms suitable for immunopurification or immunoenrichment, e.g. in which the monoclonal antibody is immobilised on a suitable solid phase such as polymer particles.

Hybridomas and other cell lines, including transformed host cell lines, which produce monoclonal antibodies according to the invention are also included in the invention.

Immunological diagnostic kits containing as specific reagent a monoclonal antibody according to the invention for the detection of MDR cells or human Pgp are also included in the invention. Such kits typically comprise a suitably labelled monoclonal antibody according to the invention, e.g. a fluorescently labelled antibody

The invention also includes the use of a monoclonal antibody according to the invention for the identification or purification of cells expressing human Pgp present in heterogeneous cell populations.

Moreover the invention also includes use of a monoclonal antibody of the invention as a therapeutic agent, e.g. to reverse MDR, and also to therapeutic compositions comprising the monoclonal antibodies. The monoclonals may be used in any of those situations in which MDR presents a clinical problem, including cancer treatment. The therapeutic compositions typically comprise an antibody according to the invention in combination with a pharmaceutically acceptable carrier or excipient.

The present invention is now described by way of illustration only in the following Examples which refer to the accompanying Figure which is a graph showing FACS analysis results for human Pgp expression on live CEM cells and its MDR variants as determined using monoclonal antibody MM4.17 (1=drug-sensitive CEM line; 2=CEM-VRL20; 3=CEM-VRL40; 4=CEM-VRL80; 5=CEM-VRL100. Abscissae: fluorescence intensity; ordinates: number of cells).

EXAMPLES

Example 1

Immunization of mice and hybridization of splenocytes with cells of the myeloma line Sp2-01/Ag.14 (ATCC CRL 1581).

BALB/c mice are immunized with an MDR variant of a human T-lymphoblastoid line already characterized and known as CEM-VBL100 (11). The derivation of the CEM-VBL100 cell line from the parent CEM cell line (ATCC CCL 119) is described by Beck et al. (11) from whom this cell line and other MDR derivatives of CEM are available. The cells of the CEM-VBL100 line proliferate actively in the presence of 100 ng/ml vinblastine or relatively high concentrations of a wide range of cytostatic agents. In order to obtain high affinity monoclonals, immunisation procedures substantially as described by Cianfriglia et al. (12) are used, with appropriate modifications. The immunization procedure involves the intraperitoneal administration of 1×10^7 CEM-VBL100 cells resuspended in PBS, live and with no pretreatment, every 15 days for 10 months. The last immunization takes place 3 days before the somatic fusion of splenocytes taken from the treated mice and involves the administration of 2×10^7 cells (resuspended in PBS) into the caudal vein and of the same number of cells intraperitoneally.

After thorough washing with serum-free culture medium, 25×10^6 splenocytes are placed in a test tube with 10×10^6 Sp2-01/Ag.14 myeloma cells (ATCC CRL 1581), pre-treated in the same way as the lymphocytes.

The mixture of spleen cells and myeloma cells is then processed as described by Cianfriglia et al. (12). Briefly, the heterokaryons obtained are distributed on microtitration plates (Costar) to obtain, after selection in HAT culture medium (H=hypoxanthine, A=aminopterin, T=thymidine, Gibco), clonal hybridomas. The culture supernatants from the cells in which increased cell growth is observed are tested for indirect immunofluorescence with the CEM drug-sensitive parent line (ATCC

CCL 119) and, simultaneously, with the CEM-VBL100 variant, to identify hybridomas secreting immunoglobulins that react selectively to the MDR immunizing cell and do not cross-react with the drug-sensitive parent line. From the hybridoma lines thus selected a line is identified, known as MM4.17, that secretes immunoglobulins of isotype IgG2a,k, which are highly reactive to the CEM-VBL100 cells, as demonstrated by an intense immunofluorescence signal, determined by flow cytometry studies, shown in Fig. 1. Flow cytometry determinations are performed using standard procedures with FITC-conjugated F(Ab')₂ anti-mouse IgG (Cappel, West Chester, PA, USA). After staining cells were fixed with 1% formaldehyde in PBS, pH7.2 and analysed on a bench-top flow cytometer (FACScan, Becton Dickinson) equipped with a 15-nW argon ion laser emitting light at a fixed wavelength of 488 nm.

The MM4.17 hybridoma is seeded onto 96-cell microtitration plates at a concentration of 0.5 cells per cell on a single layer of cells to provide nutrients (feeder layer) obtained by peritoneal lavage of BALB/c mice. Culture medium containing HT (100 µL) is added to each cell and, after 10-14 days culture, the clones are sub-cloned by the same procedure as described above a further two times. The sub-clones secrete an antibody with the same characteristics as that produced by the non-cloned line.

The MM4.17 hybridoma is cultured in vitro to obtain large quantities of monoclonal antibody from the supernatant and is injected into syngeneic BALB/c mice that have been pre-treated with Freund's incomplete adjuvant to promote the formation of ascites.

Any similar MDR human cell line may be used in place of the CEM-VBL100 for immunisation of the mice and selection of hybridoma cell lines.

Example 2

Specificity of the MM4.17 monoclonal antibody for human MDR1- P-glycoprotein

The supernatant of the hybridoma cell line MM166.4.17 (now known as MM4.17) binds only with the human MDR variants of CEM cells (as described in references 10 and 11) and is completely unreactive with its drug-sensitive parental cell line. After two cloning cycles, a cloned hybridoma cell line secreting an IgG2a, k, monoclonal immunoglobulin is isolated, characterised, and a hybridoma cell bank is set up from large-scale mycoplasma-free cultures. From this cell bank, vial samples are thawed and hybridoma cells tested for stability, sterility and specificity. Additional working cell banks are made and consistent and homogeneous batches of monoclonal antibody in either purified or crude form are obtained from the supernatants thereof. Figure 1 shows the fluorescence profiles of the monoclonal MM4.17 on CEM cells and MDR variants thereof possessing various levels of relative resistance (RR). These data clearly show the capability of MM4.17 to recognise very low RR levels (see references 10 and 13) and variations of P-glycoprotein expression on MDR cells. The MM4.17 monoclonal strongly stains only the MDR variants of the various human cell lines tested and does not significantly stain either drug-sensitive or drug-resistant cells of rodent origin. The specificity of the MM4.17 antibody is also confirmed by somatic cell genetics studies. Interspecific hybrids obtained by somatic fusion of a rodent drug-sensitive cell line with the CEM-VBL100 cells segregate according to MDR phenotype, the MDR1 gene and antigenic determinant being recognised by the MM4.17 antibody.

Example 3

Identification of the epitope of glycoprotein P recognised by the MM4.17 antibody

Peptides are synthesized that have the amino acid sequences deduced from the portions of the cDNAs that code for the extracellularly-located domains of human or murine glycoprotein

P (2, 3, 4). These peptides are incubated with the culture supernatant of the hybridoma secreting the MM4.17 antibody. An ELISA assay is used to show that the peptide corresponding to residues 739-754 of the fourth extracellular loop of Pgp (SEQUENCE ID NO. 1) is capable of reacting with the MM4.17 antibody. The other peptides from the predicted extracellular P-glycoprotein domains do not interact with antibody. Also no detectable binding is seen with mouse peptides which correspond to equivalent parts of the predicted mouse mdr1 extracellular domains. For example, MM4.17 does not cross-react with peptides representing stretches of the fourth extracellular loop of the mouse mdr1-P-glycoprotein, in spite of the relatively high degree of homology between the human and mouse peptide sequences.

To define the minimum sequence capable of reaction with the MM4.17 antibody an overlapping series of 70 tetra to decapeptides, contained in SEQUENCE ID NO. 1, are synthesized and subjected to an ELISA assay with the MM4.17 antibody. The series consists of 10 sets of tetra to decapeptides, the first set having a tetrapeptide corresponding to amino acid residues 1 to 4 and a decapeptide corresponding to amino acid residues 1 to 10 of SEQUENCE ID NO. 1 and the second set having a tetrapeptide corresponding to amino acid residues 2 to 5 and a decapeptide corresponding to amino acid residues 2 to 11 of SEQUENCE ID NO. 1 and so on for each successive set tetra to decapeptides. The ELISA assay results indicate that the octapeptide consisting of residues 2 to 9 inclusive of SEQUENCE ID NO.1 is the optimum minimum epitope. The loss of the first, and even more so that of the last, threonine produces a marked loss of the capacity to bind the MM4.17 antibody. Analysis of antibody binding to the various peptides clearly shows that: (1) binding is significantly reduced if the optimum minimum sequence is elongated in either direction; (2) the hexapeptide consisting of residues 4 to 9 inclusive of SEQUENCE ID NO.1 represents the core of the antibody recognition site, since it is found in all of the positive peptides; and (3) the specificity of the MM4.17 antibody is underlined by very significant binding values obtained with

shorter peptides, such as those consisting of residues 3 to 8, 4 to 9 and 4 to 8 inclusive of SEQUENCE ID NO.1.

Example 4

Specificity of MM4.17 for the human MDR1 gene product

The mdr specificity of the MM4.17 monoclonal is determined by assaying its binding to octapeptides of the predicted amino acid sequences of other mammalian P-glycoprotein genes which are selected for their homology to the specific epitope-forming amino acid sequence of the optimal-minimal peptide. The MM4.17 monoclonal does not bind either to the corresponding peptide (SEQUENCE ID NO. 4) of the highly divergent second member of the human mdr gene family MDR3 (identical sequences are found in mouse mdr2 and hamster pgp3 glycoproteins) or to mouse mdr1, mdr3 and hamster pgp1 and pgp2 P-glycoprotein peptides which are most similar to the MDR1 optimum minimum epitope-forming peptide .

Example 5

Identification of the nucleotide sequence and amino acid sequence of the variable region of the heavy chain of the antibody MM4.17

The cDNA sequence coding for the variable domain and hypervariable portions thereof of the heavy chain of MM4.17 is obtained as described by Orlandi et al. (14). The cDNA and corresponding predicted amino acid sequences are given below in SEQUENCE ID NO. 2 and the amino acid sequence alone is given as SEQUENCE ID NO. 3. The Complementarity Determining Regions (CDRs) are residues 32 to 36 inclusive (for CDR1), residues 51 to 67 (for CDR2) and residues 100 to 111 (for CDR3) of the heavy chain variable domain amino acid sequence given in SEQUENCE ID NOS 2 and 3.

Example 6Comparison of the binding affinity of MM4.17 for Pgp on MDR and drug-sensitive cells with previous monoclonals

The MM4.17 and previous monoclonals C219, C494, JSB1, MRK-16 and MC57 are compared for binding to CEM, CEM-VBL10 and CEM-VBL100 in flow cytometry experiments in which the staining characteristics of fluorescently labelled versions of the antibodies are measured. The results obtained are given in the Table below. The monoclonals were used in purified form and at the same protein concentration (10 μ g/ml). The results clearly show that the MM4.17 has substantially higher binding affinity for MDR cell lines and is the monoclonal which unequivocally binds to the CEM-VBL10 cell line.

Table

Mab	Pgp domain	cells		CEM	CEM-VBL10	CEM-VBL100	CEM/CEM-VBL10		CEM-VBL100	
							fluorescent profile		fluorescent profile	
C219	cyt ¹	+ ⁻²	+ ⁻			+	yes ³			yes
C494	cyt	+ ⁻	+ ⁻			++	yes			yes
JSB1	cyt	+ ⁻	+ ⁻			+	yes			yes
MRK-16	ext ⁴	-	+ ⁻			+++	yes			no
MC57	ext	-	+ ⁻			+++	yes			no
MM4.17	ext	-	+++			++++	no			no

4 In the above Table the reference numbers have the following significances:

- 1 - cytoplasmic epitope recognition (the cells have to be fixed or permeabilised for Mab staining.
- 2 - level of reactivity: -, negative; +⁻, from 5 to 30%; +, from > 30 to 60%; ++, from > 60 to 90%; +++, from > 90 to 100%; ++++, 100% of the cells stained but with a higher fluorescent intensity.
- 3 - the fluorescent profiles of the cells show an overlapping area of reactivity.
- 4 - extracellular epitope recognition in live human MDR cells.

REFERENCES

1. Kohler G. and Milstein C., Nature, 256, 495, 1975
2. Ueda et al. (1987), PNAS, 84, 3004-3008.
3. Shen et al. (1986), Mol. Cell. Biol., 6, 4039-4044.
4. Chen et al. (1987), Cell., 47, 381-389.
5. Cenciarelli et al., (1991) Int. J. Cancer, 47, 533-544, .
6. Arceci et al. (1993) Cancer Res., 53, 310-317.
7. Mechtner & Roninson (1992) PNAS, 89, 5824-5828.
8. Georges et al. (1993) JBC, 268, 1792-1798.
9. Hamada and Tsuruo (1986), PNAS, 83, 7785-7789.
10. Cianfriglia et al. (1990) Int. J. of Cancer, 45, 95-103.
11. Beck et al. (1979) Cancer Res., 39, 2070-2077.
12. Cianfriglia et al., (1986) Meth. in Enzymol., 121, 193-210,
13. Wyler et al. (1993) Proceedings of the 84th Annual Meeting of AACR, Abs. No. 1837, page 309.
14. Orlandi et al., (1989) PNAS, 86, 3833-3837 .
15. Kartner et al. (1985), Nature, 316, 820-823.
16. Scheper et al. (1988), Int. J. Cancer, 42, 389-394.

SEQUENCE LISTING

(1) GENERAL INFORMATION:

(i) APPLICANT:

- (A) NAME: Istituto Superiore di Sanita
- (B) STREET: Viale Regina Elena 299
- (C) CITY: 00161 Rome
- (E) COUNTRY: Italy
- (F) POSTAL CODE (ZIP): 00161
- (G) TELEPHONE: 003964444990
- (H) TELEFAX: 003964440067
- (I) TELEX: RM610071

(ii) TITLE OF INVENTION: Monoclonal Antibodies to glycoprotein P

(iii) NUMBER OF SEQUENCES: 4

(iv) COMPUTER READABLE FORM:

- (A) MEDIUM TYPE: Floppy disk
- (B) COMPUTER: IBM PC compatible
- (C) OPERATING SYSTEM: PC-DOS/MS-DOS
- (D) SOFTWARE: PatentIn Release #1.0, Version #1.25

(EPO)

(v) CURRENT APPLICATION DATA:

APPLICATION NUMBER: WO PCT/EP93/

(2) INFORMATION FOR SEQ ID NO: 1:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 16 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO

(v) FRAGMENT TYPE: internal

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 1:

Phe	Thr	Arg	Ile	Asp	Asp	Pro	Glu	Thr	Lys	Arg	Gln	Asn	Ser	Asn	Leu
1				5					10					15	

(2) INFORMATION FOR SEQ ID NO: 2:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 369 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iii) ANTI-SENSE: NO

(v) FRAGMENT TYPE: internal

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Mus musculus

(B) STRAIN: BALB/c

(ix) FEATURE:

(A) NAME/KEY: CDS

(B) LOCATION: 1..369

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 2:

CAG GTC CAA CTG CAG GAG TCT GGA GGA GAC TTA GTG AAG GAT CCT GGA	48
Gln Val Gln Leu Gln Glu Ser Gly Gly Asp Leu Val Lys Asp Pro Gly	
1 5 10 15	
GGG TCC CTG AAA CTC TCC TGT GCA GCC TCT GGA TTC ACT TTC AGT AGA	96
Gly Ser Leu Lys Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Arg	
20 25 30	
TAT GGC ATG TCT TGG GTT CGC CAG ACT CCA GAC AAG AGG CTG GAG TGG	144
Tyr Gly Met Ser Trp Val Arg Gln Thr Pro Asp Lys Arg Leu Glu Trp	
35 40 45	
GTC GCA ACC ATT AGT AGC GGT GGT AGT TAC ACC TAC TTT CCA GAC AGT	192
Val Ala Thr Ile Ser Ser Gly Gly Ser Tyr Thr Tyr Phe Pro Asp Ser	
50 55 60	
GTG AAG GGG CGA TTC ACC ATC TCC AGA GAC AAT GCC AAG AAC ACC CTG	240
Val Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Asn Thr Leu	
65 70 75 80	
TAC CTG CAA GTG AGC AGT CTG AAG TCT GAG GAC ACA GCC ATG TAT TAC	288
Tyr Leu Gln Val Ser Ser Leu Lys Ser Glu Asp Thr Ala Met Tyr Tyr	
85 90 95	
TGT GCA AGA CCT GCG GAA TTT AGG GGT TAC TCC TGG TTT GCT TAC TGG	336
Cys Ala Arg Pro Ala Glu Phe Arg Gly Tyr Ser Trp Phe Ala Tyr Trp	
100 105 110	
GGC CAA GGG ACC ACG GTC ACC GTC TCC TCA GGT	369
Gly Gln Gly Thr Thr Val Thr Val Ser Ser Gly	
115 120	

(2) INFORMATION FOR SEQ ID NO: 3:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 123 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 3:

Gln	Val	Gln	Leu	Gln	Glu	Ser	Gly	Gly	Asp	Leu	Val	Lys	Asp	Pro	Gly
1				5					10					15	
Gly	Ser	Leu	Lys	Leu	Ser	Cys	Ala	Ala	Ser	Gly	Phe	Thr	Phe	Ser	Arg
			20					25					30		
Tyr	Gly	Met	Ser	Trp	Val	Arg	Gln	Thr	Pro	Asp	Lys	Arg	Leu	Glu	Trp
		35					40					45			
Val	Ala	Thr	Ile	Ser	Ser	Gly	Gly	Ser	Tyr	Thr	Tyr	Phe	Pro	Asp	Ser
		50				55					60				
Val	Lys	Gly	Arg	Phe	Thr	Ile	Ser	Arg	Asp	Asn	Ala	Lys	Asn	Thr	Leu
65					70				75					80	
Tyr	Leu	Gln	Val	Ser	Ser	Leu	Lys	Ser	Glu	Asp	Thr	Ala	Met	Tyr	Tyr
				85					90					95	
Cys	Ala	Arg	Pro	Ala	Glu	Phe	Arg	Gly	Tyr	Ser	Trp	Phe	Ala	Tyr	Trp
			100					105					110		
Gly	Gln	Gly	Thr	Thr	Val	Thr	Val	Ser	Ser	Gly					
		115					120								

(2) INFORMATION FOR SEQ ID NO: 4:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 8 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO

(v) FRAGMENT TYPE: internal

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo sapiens

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 4:

Gly Pro Gly Asp Asp Ala Val Lys

1

5

Claims


1. A monoclonal antibody that recognise a structurally continuous and extracellularly-located epitope of human Pgp consisting of a continuous epitope-forming amino acid sequence.
2. A monoclonal antibody according to Claim 1 in which the said epitope-forming amino acid sequence is located on the fourth extracellular loop of human Pgp.
3. A monoclonal antibody according to Claim 2 in which the said epitope-forming amino acid sequence is contained in the peptide of SEQUENCE ID NO. 1.
4. A monoclonal antibody according to Claim 3 in which the said epitope-forming amino acid sequence comprises at least 5 contiguous amino acid residues of the 8 amino acid sequence consisting of residues 2 to 9 inclusive of SEQUENCE ID NO. 1.
5. A monoclonal antibody according to Claim 4 in which the 5 amino acids are residues 3 to 7 inclusive of SEQUENCE ID NO. 1.
6. A monoclonal antibody to human Pgp comprising a variable domain having CDRs which are the same as or variants of the CDRs of the heavy chain variable domain amino acid sequence of SEQUENCE ID NOS. 2 or 3.
7. A monoclonal antibody according to Claim 6 in which each CDR is at least 70% homologous to the corresponding CDR of SEQUENCE ID NOS. 2 or 3.
8. A monoclonal antibody according to any of the preceding

Claims which has a binding affinity for Pgp similar to that of the MM4.17 monoclonal antibody.

9. A monoclonal antibody according to any of the preceding Claims which is fluorescently labelled or immobilised on a solid phase.
10. A method for the preparation of a monoclonal antibody according to any of the preceding claims comprising somatic cell fusion of spleen cells obtained from an animal which has been immunised with human MDR cells, or peptides corresponding to extracellular domains of human Pgp or fragments thereof.
11. A method according to Claim 10 in which a peptide corresponding to a structurally continuous extracellular domain of human Pgp or a part thereof is used to screen for the desired monoclonal antibody.
12. Hybridoma and transformed host cell lines producing a monoclonal antibody according to any of claims 1-8.
13. An immunological diagnostic kit containing as specific reagent a monoclonal antibody according to any of Claims 1 to 9 for the detection of human MDR cells or human Pgp.
14. Use of a monoclonal antibody according to any of Claims 1 to 9 for the identification or purification of cells expressing human Pgp present in heterogeneous cell populations.
15. Use of a monoclonal antibody according to any of Claims 1 to 8 as a therapeutic agent to reverse MDR.
16. Use of a monoclonal antibody according to any of Claims 1 to 8 to reverse MDR.

17. A therapeutic composition comprising a monoclonal antibody according to any of Claims 1 to 8 in combination with a pharmaceutically acceptable carrier or excipient.

FIGURE



INTERNATIONAL SEARCH REPORT

PCT/EP 93/01533

International Application No

I. CLASSIFICATION OF SUBJECT MATTER (if several classification symbols apply, indicate all) ⁶		
According to International Patent Classification (IPC) or to both National Classification and IPC		
Int.Cl. 5 C12P21/08; C12N5/20; G01N33/574; G01N33/577 A61K39/395		
II. FIELDS SEARCHED		
Minimum Documentation Searched ⁷		
Classification System	Classification Symbols	
Int.Cl. 5	C12P ; C12N ; G01N ; A61K	
Documentation Searched other than Minimum Documentation to the Extent that such Documents are Included in the Fields Searched ⁸		
III. DOCUMENTS CONSIDERED TO BE RELEVANT⁹		
Category ¹⁰	Citation of Document, ¹¹ with indication, where appropriate, of the relevant passages ¹²	Relevant to Claim No. ¹³
A	JOURNAL OF CELLULAR PHARMACOLOGY vol. 2, no. 3, 1991, BERLIN, GERMANY pages 165 - 170 M. CHEVALLIER-MULTON ET AL. 'Importance of the fourth external loop in immunogenicity of the multidrug resistance related P glycoprotein.' see the whole document ----	1-17
P,X	PROCEEDINGS OF THE AMERICAN ASSOCIATION FOR CANCER RESEARCH, ANNUAL MEETING. vol. 34, no. 0, March 1993, page 317 M. CIANFRIGLIA ET AL. 'Precise mapping of a continuous human-specific epitope in the fourth loop of the P glycoprotein.' see abstract 1889 -----	1-17
<p>¹⁰ Special categories of cited documents : ¹⁰</p> <p>"A" document defining the general state of the art which is not considered to be of particular relevance</p> <p>"E" earlier document but published on or after the international filing date</p> <p>"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>"O" document referring to an oral disclosure, use, exhibition or other means</p> <p>"P" document published prior to the international filing date but later than the priority date claimed</p> <p>"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step</p> <p>"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.</p> <p>"&" document member of the same patent family</p>		
IV. CERTIFICATION		
Date of the Actual Completion of the International Search	Date of Mailing of this International Search Report	
25 AUGUST 1993	23 -09- 1993	
International Searching Authority	Signature of Authorized Officer	
EUROPEAN PATENT OFFICE	NOOIJ F.J.M.	

INTERNATIONAL SEARCH REPORT

Int l application No.

PCT/EP 93/01533

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
Remark : Although claims 15 & 16 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.
2. ☐ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.